

HEALTH CARE PLAN FOR A CHILD/YOUNG PERSON WITH EPILEPSY

Photo

PERSONAL DETAILS

Student Full Name:

Tutor Group:				
Date of Birth				
Date Care Plan Completed:				
CONTACT INFORMATION				
Family Contact 1	Family Contact 2			
Name:	Name:			
Phone No: (Home)	Phone No: (Home)			
(Work):	(Work):			
(Mobile):	(Mobile):			
Relationship:	Relationship:			
GP Name:	Consultant Name:			
GP Address:	Hospital/Clinic:			
Telephone Number:	Telephone Number:			
Please complete the form below with as much de	etail as possible.			
Type of Seizure PARTIAL SEIZURES				
 □ Remains conscious □ With motor symptoms jerking and stiffening □ With Touch, smell, hearing, taste, and sight symptoms □ With autonomic symptoms heart rate change, internal sensations □ With psychic symptoms in a dream like state □ Complex partial seizures Conscious but impaired movements of the mouth and face (e.g., lip smacking, chewing, and swallowing movements), the hands and arms (e.g., fumbling, picking, and tapping movements), vocalizations (e.g., grunts or repetition of words or phrases). Absence Seizures (Formerly called petit mal) 				
 □ Typical brief - episodes of staring, blinking, unaware of surroundings; usually lasts less than 10 seconds but may last up to 20 seconds □ Atypical - staring spells lasting between 5 to 30 seconds, eye blinking or slight jerking movement of the lips may occur; partial reduction in responsiveness. □ Myoclonic - brief jerks of a muscle or group of muscles; usually involving the neck, shoulders, and upper arms. □ Atonic Sudden loss of muscle strength, eyelids may droop, head may nod, objects may be dropped, or the child may fall to the ground; usually lasts less than 15 seconds, injury is common. □ Clonic Rhythmic- jerking movements of the arms and legs, maybe generalised. □ Tonic- sudden stiffening movements of the body, arms, or legs involving both sides of the body; usually last less than 20 seconds. Tonic-Clonic (Formerly called grand mal) □ Convulsive seizures, body briefly stiffens followed by a jerking motion of the arms and legs; loss of consciousness and falls frequently occur, excessive saliva production may be present, possible loss of bowel and bladder control; usually lasts a couple of minutes, the child is often tired or confused after the seizure and may want to go to sleep. 				

If your son/daughter experiences any other symptoms before, during or after a seizure, please provide more information below:							
Other symptoms							
(Specify:)							
Advice							
Known triggers – if any:							
Date of last seizure:							
Usual duration of seizure: Recovery time:							
Actions – Please advise us of how you care for your son/daughter both during and after a seizure.							
Current medication sum		T 5	F	0' 1- E((((1)'-			
Anti-Epileptic Medication Taken	Method Oral, Rectal, Gel	Dose (mg or ml)	Frequency (times daily)	Side Effects of this medication			
	(If rectal Diazepam or Buccal Midazolam are being used then you will be sent a separate form to fill in for this)		, , , ,				
SPECIAL CONSIDERATIONS: (discuss further with Student Services/Health & Safety)							
Swimming:							
Physical Education:							
Practical lessons:							
School Trips:			(la a a la a la la la la la la la la la l				
Any other information you feel may be important for the school to be aware of?							
treatment/medication of CONFIDENTIALITY : F	ninded of the importance or ongoing concerns/chang	ges in episode apid access, t	patterns.				
Signature of Parent/Guardian:	board in the staff room and in subject department offices. Signature of Parent/Guardian: Print Name: Date:						
Office Use Only							
Health Care Plan Agreed by:	re Plan Agreed by: Copy sent to parent date:						
Date Care Plan Created:	Added to Medical Register on date:						